Turnitin Artikel

by Nyoman Suci Widyastiti

Submission date: 01-Sep-2025 10:00AM (UTC+0700)
Submission ID: 2739201334
File name: Submit_Format_manuscript_IRF_Nyoman_Suci_W-Andine_DIMJ.docx (413.99K)
Word count: 3293
Character count: 20049

Diponegoro International Medical Journal

Vol.X, No.X, July/December XXXX
Available online at: https://ejournal2.undip.ac.id/index.php/dimj

Immature Reticulocyte Fraction in Thalassemia Major Receiving Single and Combination Iron Chelation Therapy

Andina Putri Aulia[1], Nyoman Suci Widyastiti[2*]

[1]*Clinical Pathology Residency Program, Faculty of Medicine, Universitas Diponegoro, Semarang / Staff of the Department of Clinical Pathology, Faculty of Medicine, Sultan Agung Islamic University, Semarang, Indonesia*
[2]*Department of Clinical Pathology, Faculty of Medicine, Universitas Diponegoro University, Semarang, Indonesia*
* Corresponding author's Email: nyoman.suci@fk.undip.ac.id

Article history:

Received 20XX
Revised 20XX
Accepted 20XX
Available online 20XX

Keywords:

Thalassemia
Erythropoiesis
Iron chelation
IRF
Reticulocyte

Abstract

Background: Ineffective erythropoiesis in thalassemia major causes anemia and requires regular blood transfusions. However, routine transfusions cause iron accumulation, which can be removed through iron chelation therapy (single or combination). To monitor erythropoietic activity non-invasively, Immature Reticulocyte Fraction (IRF) and reticulocyte count have been proven effective, eliminating the need for invasive bone marrow examination.
Objective: This study aimed to determine the differences in IRF percentage and reticulocyte count between thalassemia major patients receiving single and combination iron chelation therapy.
Methods: An observational analytical study with a cross-sectional design was conducted at the Paediatric Haematology and Oncology Clinic of Dr. Kariadi General Hospital, Semarang, from November 2023 to January 2024. Subjects were thalassemia patients aged ≤ 18 years and receiving more than 10 regular transfusions. IRF and reticulocyte count were examined using fluorescence flow cytometry on whole blood samples with EDTA anticoagulant.
Results: A total of 71 thalassemia major patients were enrolled, consisting of 40 males (56%) and 31 females (44%). The mean IRF percentage was 11.62±5.4% in thalassemia major patients receiving single iron chelation therapy and 17.01±10.0% ($p<0.015$) in the group that was administered combination therapy. Mean reticulocyte count was 0.106±0.08 x106/µL in thalassemia major patients receiving single iron chelation therapy and 0.110±0.08 x106/µL in the group administered combination therapy ($p<0.83$).
Conclusion: There is a difference in IRF between thalassemia major patients, with high erythropoietic activity in the group receiving combination iron chelation therapy.

1. Introduction

Thalassemia is a hereditary hemolytic disease caused by mutations in the globin gene that disrupt globin chain synthesis. This disease is characterized by decreased or absent synthesis of the α, β, and/or other globin chains that form the normal structure of the main hemoglobin molecule in adults. Furthermore, decreased synthesis of α and β chains causes the inability of the body to form red blood cells properly, leading to anemia in childhood that persists throughout life [1,2]. Based on clinical abnormalities, thalassemia is divided into major, intermediate, and minor. Thalassemia major is the most severe form, requiring regular blood transfusions. The genetic classification is based on abnormalities in the affected globin chain subunits, specifically α-thalassemia, β-thalassemia, α-β-thalassemia, δ-β-thalassemia, and δ-γ-thalassemia. Among these types, β-thalassemia spectrum is the disease with the most mutations [1],[3].

ISSN XXXX-XXXX (online) XXXX-XXXX (print) © 20XX The Authors. Published by Universitas Diponegoro.
This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/)
doi: http://dx.doi.org/.....

As the hereditary hemolytic disease with the highest prevalence, thalassemia is a very serious health problem, accounting for the deaths of children yearly. According to data from the World Health Organization (WHO) in 2020, the global incidence of thalassemia reached approximately 54.348 million people, or 7% of the world's population. The highest incidence of 21.7 million people (40%) occurred in Asian countries. Data from 2021 showed that the global prevalence of thalassemia was estimated to be 156.74 million people, or approximately 20% of the total global population. Based on the data from the Indonesian Thalassemia Foundation, there has been a continuous increase in cases. Since 2012, there have been 4,896 cases, which have steadily increased to 10,973 as of June 2021. Among the total cases, 30% are in Central Java, ranking second nationally after West Java. In terms of financing, the 2020 BPJS Kesehatan data recorded a consistent rise in healthcare cost burden from 2014 to 2020. Thalassemia ranks fifth among non-communicable diseases, after heart disease, kidney failure, cancer, and stroke, with an estimated cost of IDR 2.78 trillion in 2020 [3–6]. Kariadi General Hospital in Semarang serves as a referral hospital for thalassemia patients, where the demand for blood supplies for therapy is extremely high.

Thalassemia major is the most severe clinical condition, which occurs because the hemoglobin-coding gene on two chromosome alleles is defective. This causes ineffective erythropoiesis, leading to anemia and the need for regular blood transfusions. Patients often require blood transfusions from the first year of development, typically between 6 and 24 months of age, which continues during the lifetime. However, regular blood transfusions (>10x) cause iron accumulation in the body, which affects several organs, suggesting the need for single or combination iron chelation therapy [7],[8]. Iron, which the body cannot naturally excrete, requires the assistance of chelators to facilitate its excretion. The three types of iron chelators currently in use are deferoxamine (DFO), deferiprone (DFP), and deferasirox (DFX). Iron chelation therapy is expected to reduce accumulation by binding iron and increasing hepcidin, thereby affecting erythropoiesis [3],[7].

Erythropoietic activity can be assessed through peripheral blood reticulocyte examination, making it easier to perform than invasive bone marrow examination. Among various markers, Immature Reticulocyte Fraction (IRF) is a sensitive early indicator for evaluating erythropoietic regeneration through hematological analysis. In thalassemia, an imbalance in the production of α and β globin chains leads to a decreased survival of erythrocytes and their precursors. Specifically, in β-thalassemia, the accumulation of excess α-chains causes oxidative stress and damage to cell membranes. This triggers apoptosis, where damaged red blood cell precursors are gradually phagocytosed by activated macrophages in the bone marrow. Ineffective erythropoiesis in thalassemia patients is characterized by an increase in erythroid precursors and progenitor cells, leading to ineffective erythropoiesis and anemia. In this context, administration of single or combination iron chelation to thalassemia major patients is performed to reduce ferritin levels, which are expected to affect the quality of erythropoiesis. Previous studies have shown that high transferrin and hepcidin levels can decrease iron bioavailability for erythroblasts, causing an increase in erythroid precursors in rats with β-thalassemia [9],[10],[11]. Therefore, iron chelation therapy is expected to improve anemia in thalassemia major patients [12,13]

Studies on the differences in the percentage of IRF and reticulocyte count between thalassemia major patients receiving single and combination iron chelation therapy are still limited. Therefore, this study aimed to provide an overview of the percentage of IRF and reticulocyte count in thalassemia major patients, offering insight into the erythropoietic status

2. Methods

This analytical observational study with a cross-sectional design was conducted from November 2023 to January 2024 at the Pediatric Haematology and Oncology Clinic of Dr. Kariadi General Hospital, Semarang. The inclusion criteria were patients aged ≤ 18 years who received more than 10 regular transfusions. The exclusion criteria were patients with a history of splenectomy and renal failure. Data, including age, gender, and laboratory

Published by Universitas Diponegoro.
doi: http://dx.doi.org/.....

parameters, were collected from the electronic medical records (EMRs) of Dr. Kariadi General Hospital, Semarang. A total sample of 245 patients was obtained, where 174 were excluded due to incomplete data, and 71 patients were included in the study sample.

Data were obtained through electronic medical records. Haemoglobin levels, erythrocyte count, IRF percentage, and reticulocyte count were measured at Dr. Kariadi General Hospital using a Sysmex XN-1000 haematology analyzer. In comparison, ferritin was measured using patients' serum through the Vidas Enzyme-Linked Fluorescent Assay (ELFA) method.

Data were processed using SPSS 23, and a normality test was conducted using the Kolmogorov-Smirnov test. Analysis of differences was performed using the independent t-test and the Mann-Whitney test. Statistical test results were considered significant when the p-value was <0.05.

3. Results

The data were obtained from 71 patients who were divided into two groups, namely single and combination iron chelation therapy, as shown in Table 1. The prevalence of disease in this study was higher in males than in females, with a median age of 11 years (range, 4-17). All patients received oral iron chelation therapy. In the single iron chelation therapy group, 32 (72.7%) patients received Deferasirox (DFX) and 12 (27.2%) were given Deferiprone (DFP). Most patients received transfusions every 4 weeks (45%).

Table 1. Characteristics of Patient's Data

Variables	**Total n = 71**	**Iron Chelation Therapy**	
		Single n = 44 (62%)	Combination n = 27 (38%)
Male	40 (56.3%)	21(47.7%)	19 (70.3%)
Female	31 (43.6%)	23 (52.2%)	8 (29.6%)
Age#	11 (4-17)	10 (5-16)	13 (4-17)
Hemoglobin# (g/dL)	7.46 (4-10.1)	7.36 (4-10.1)	7.62 (5-9.7)
Erythrocytes# (106 /uL)	3.27 (1.61-4.25)	3.24 ± 0.52	3.32 ± 0.50
Iron chelation			
Deferasirox		32 (72.7%)	
Deferipron		12 (27.2%)	
Transfusion frequency			
Every 2 weeks	23 (31.9%)	12 (27.2%)	11 (40.7%)
Every 3 weeks	15 (20.8%)	7 (15.9%)	8 (29.6%)
Every 4 weeks	33 (45.8%)	25 (56.8%)	8 (29.6%)
Ferritin# (µg/dL)	3262 (338–13999)	2678.5 (338-6760)	4212 (976-13999)

SD, standard deviation; min, minimum; max, maximum; #, non-normal data distribution

Data were analyzed for normality using the Kolmogorov-Smirnov test. Analysis continued with a test for differences in IRF percentage, reticulocyte count, and ferritin levels between the two groups. The values obtained were p = 0.015, p = 0.83, and p = 0.024, respectively, as shown in Table 2.

In this study, a significant difference was observed between the IRF percentage in single and combination iron chelation therapy groups. The mean reticulocyte count in the combination was descriptively higher than in the single iron chelation therapy group. However, there was no significant difference between the two groups.

Table 2. Difference test of IRF percentage and reticulocyte count in thalassemia major patients with single and combination iron chelation therapy

Variables	**Total n = 71**	**Iron Chelation Therapy**		**p**
		Single n = 44 (62%)	Combi-nation n = 27 (38%)	
IRF (%)^	13.6 ± 7.92	11.62 ± 5.4	17.01 ± 10.0	0.015*
Reticulocyte (106/uL)^	0.107 ± 0.07	0.106 ± 0.08	0.110 ±0.08	0.83

§Mann-Whitney test; ^Independent t-test ; *p< 0,05

4. Discussion

This study aimed to determine the differences in IRF percentage and reticulocyte count between pediatric thalassemia major patients receiving single and combination iron chelation therapy. The majority

Published by Universitas Diponegoro.
doi: http://dx.doi.org/.....

of subjects were male (56.3%) and aged over 5 years (94.3%). Based on age, these results were in line with data from a previous study showing that the average age of thalassemia patients was 11.1±3.71 years.[14] However, this result was in contradiction with previous studies that reported a higher number of female patients compared to male patients in thalassemia.[15] According to data from the Indonesian Thalassemia Foundation (FKHTI) in 2014, the number of patients was 6,439, with 3,219 males and 3,220 females [16].

A total of 44 patients (62%) received single iron chelation therapy, with 32 given DFX and 12 had DFP. In line with the results, approximately 31.9%, 20.58%, and 45% of patients received transfusions every 2, 3, and 4 weeks. Similarly, a previous study reported that thalassemia patients started transfusions at an average age of 3.78 years, with a frequency of approximately once a month (87.5%) [17]. Median ferritin levels were higher in the group receiving combination iron chelation therapy compared to the single. These results varied compared to another report, where ferritin levels in combination with iron chelation therapy were significantly lower than in single therapy [7]. The variation was because the determination of ferritin levels in this study was conducted without considering the duration of therapy and previous levels.

The results found a significant difference in IRF percentage between thalassemia major patients receiving single and combination iron chelation therapy. The IRF percentage was obtained from the sum of the percentages of Medium-fluorescence reticulocyte (MFR) or semi-mature reticulocyte and High-fluorescence reticulocyte (HFR) or immature reticulocytes. Meanwhile, the percentage of Low-fluorescence reticulocyte (LFR) or mature reticulocyte was obtained by subtracting 100% from the IRF percentage. This suggested that a higher IRF percentage correlated with a lower LFR. An increase in IRF represents a rise in immature reticulocytes that have recently been released from the bone marrow. This shows an erythropoietic response from the bone marrow, characterized by the accelerated release of immature reticulocytes due to anemia. Therefore, IRF is a better indicator of bone marrow function compared to other reticulocyte parameters [13]. Andriyani et al. (2021) showed that the IRF percentage in transfusion-dependent thalassemia patients with severe anemia was significantly higher than in moderate cases. Furthermore, thalassemia major patients showed low hepcidin expression and high ferroportin expression, causing enhanced iron absorption through the gastrointestinal tract. Transfusion iron overload and increased gastrointestinal iron absorption are causes of secondary hemosiderosis in thalassemia patients [1],[18].

Chelation therapy aims to eliminate and prevent iron overload by clearing excess iron and maintaining natural levels. Therefore, the administration of iron chelation can reduce the toxic effects of iron overload. Based on the mechanism of operation, iron chelation enters cells, binds free iron, and removes it from the body [19]. The combination of DFP and DFX is more effective in clearing the heart and liver of iron siderosis. This is due to the synergistic effect as well as the ability of DFX and DFP to enter cells and chelate intracellular iron [20]. Karami et al. (2017) showed a significant difference in liver iron concentration (LIC) in β-thalassemia major patients before and after combination iron chelation therapy (DFP and DFX) [20].

Iron restriction with combination therapy can reduce TSat and erythroid iron intake, thereby limiting heme synthesis. This shows the potential to limit hemichrome production, which hinders the toxicity associated with excess α-globin chains. Reducing hemichrome formation can also enhance the quality of the RBCs produced, thereby increasing lifespan and the number of circulating RBCs. Additional beneficial effects of iron restriction include low hypoxia and EPO synthesis, with increased extramedullary erythropoiesis and the erythroid progenitor/RBC ratio [12]. Deferiprone (DFP) increases hepcidin expression, altering iron distribution by moving it out of parenchymal cells and into circulating transferrin, enhancing iron-mediated signaling to hepcidin expression. Specifically, DFP binds iron when there is an overload and removes excess from the body. It also has antioxidant effects, preventing free radicals, ferroptosis, and cuproptosis, thereby inhibiting tissue

Published by Universitas Diponegoro.
doi: http://dx.doi.org/.....

damage caused by excess ferritin accumulation [21]. In comparison, DFX binds iron at a 2:1 ratio, reducing toxicity by binding trivalent iron with high affinity to a stable form that is excreted in the urine [22].

In this study, reticulocyte count in the combination iron chelation therapy group was descriptively higher. There was a statistically significant difference between the two groups. This was because reticulocyte count represented the total number of reticulocytes, while IRF percentage was calculated by summing the MFR and HFR percentage. The principle of reticulocyte examination is flow cytometry. RNA/DNA in reticulocyte is stained, then the stained cells are detected and counted based on a two-dimensional distribution, namely, forward scattered light and side fluorescent light [1],[23].

A limitation of this study is the inability to consider the duration of iron chelation therapy as well as ferritin levels before and after iron chelation. Therefore, the amount of iron released to influence ineffective erythropoiesis in thalassemia major patients is not determined during the analysis.

5. Conclusions and Suggestions

This study shows that there is a significant difference in IRF percentage among thalassemia major patients receiving single and combination iron chelation therapy ($p<0.001$). However, reticulocyte count does not show a significant difference between the two groups. This suggests that erythropoietic activity in thalassemia major patients receiving combination iron chelation therapy is higher. Moreover, further studies are needed to periodically determine IRF percentage and reticulocyte count in thalassemia major patients receiving single and combination iron chelation therapy, as well as to measure ferritin levels before and after administration.

Ethical Approval

The ethical approval was received from the Research Ethics Committee of Dr. Kariadi General Hospital, Semarang, under number 1697/EC/KEPK-RSDK/2024.

Conflicts of Interest

The authors declare no conflict of interest.

Funding

This research was conducted as part of a study supported by the Faculty of Medicine, Universitas Diponegoro, research grant no. 572/UN7.F4/PP/VI/2024.

Author Contributions

APA: conceptualization, methodology, formal analysis, investigation, resources, data curation, writing-original draft, visualization. NSW: conceptualization, methodology, formal analysis, investigation, resources, data curation, writing-original draft, visualization, supervision, project administration, funding acquisition.

Acknowledgments

I would like to express my sincere gratitude to the Faculty of Medicine, Universitas Diponegoro, for their support and facilitation throughout this research. Special thanks are also extended to RSUP Dr. Kariadi for providing the facilities and assistance essential for the data collection and successful completion of this study. I am grateful to all the staff and colleagues who contributed valuable insights and encouragement during the research process.

References

1. Andriyani R, Prihatni D, Parwati I, Rostini T. Difference in Immature Reticulocyte Fraction Percentage between Moderate and Severe Anemia in Transfusion-Dependent Thalassemia. Majalah Kedokteran Bandung. 2021;53(4):235–9.
2. Canis Paloma IDAN. Talasemia : sebuah Tinjauan Pustaka. Biocity Journal of Pharmacy Bioscience and Clinical Community. 2023;1(2):89–100.
3. Abarca RM. Talasemia Genetik Dasar dan Pengelolaan Terkini. Nuevos sistemas de comunicación e información. 2021. 2013–2015 p.
4. Nurvinanda. SuhanaHKusumajayaR. Faktor Faktor Yang Berhubungan Dengan Kejadian Thalassemia Beta Mayor Pada Anak. Jurnal Penelitian Perawat Profesional. 2023;5(November):1713–24.
5. Amalia Y. Difference of Het Re Level In Thalassemia β Minor And Iron Deficiency Anemia. Vol. 9, STRADA Jurnal Ilmiah Kesehatan. 2020. p. 1635–41.
6. Widyawati. Talasemia Penyakit Keturunan, Hindari dengan Deteksi Dini [Internet]. Kemenkes RI. 2022. Available from:

https://sehatnegeriku.kemkes.go.id/baca/rilis-media/20220510/5739792/talasemia-penyakit-keturunan-hindari-dengan-deteksi-dini/

7. Jhinger P, Sobti PC, Kaushal S, Kakkar S. Combination of two oral iron chelators in patients with thalassemia major. Pediatric Hematology Oncology Journal. 2018;3(3):55–8.
8. Gao C, Li L, Chen B, Song H, Cheng J, Zhang X, et al. Clinical outcomes of transfusion-associated iron overload in patients with refractory chronic anemia. Patient Prefer Adherence. 2014;8:513–7.
9. Anovadia DrB, Trivedi DrDP, Savjiyani DrD, Popat DrVC. Clinical significance of immature reticulocyte fraction (IRF) and reticulocyte maturity indices in differential diagnosis of anemia. International Journal of Clinical and Diagnostic Pathology. 2023;6(1):16–9.
10. Saad HKM, Rahman AAA, Ghani ASA, Wan Taib WR, Ismail I, Johan MF, et al. Activation of STAT and SMAD Signaling Induces Hepcidin Re-Expression as a Therapeutic Target for-Thalassemia Patients. Vol. 10, Biomedicines. 2022.
11. Rivella S. Ineffective erythropoiesis and thalassemias Stefano. Curr Opin Hematol. 2019;16(3):187–94.
12. Rivella S. Iron metabolism under conditions of ineffective erythropoiesis in β-Thalassemia. Blood. 2019;133(1):51–8.
13. Adane T, Asrie F GZ. Clinical Utility of Immature Reticulocyte Fraction. J Clin Chem Lab Med. 2021;4(9):1–5.
14. Widodo S, Widyastiti NS, Limijadi EKS, Hendrianingtyas M, Retnoningrum D, Ariosta A, et al. Insulin-like Growth Factor-1 and Calcium Ion Levels are Negatively Associated with Serum β-Cross Laps in Multi-transfused β-thalassemia Major Patients. The Indonesian Biomedical Journal [Internet]. 2025 Aug 25;17(4):382–8. Available from: https://inabj.org/index.php/ibj/article/view/3706
15. Caroline POL, Widyastiti NS, Ariosta, Pratiwi R, Retnoningrum D, Ngestiningsih D, et al. The differences of 25-hydroxyvitamin D and malondialdehyde levels among thalassemia major and non-thalassemia. Bali Medical Journal. 2021 Aug 1;10(2):617–22.
16. Keputusan Menteri Kesehatan Republik Indonesia Nomor HK. 01.07/ MENKES/ 1/2018. PEDOMAN NASIONAL PELAYANAN KEDOKTERAN TATA LAKSANA THALASEMIA. 2018.
17. Sawitri H, Husna CA. Karakteristik Pasien Thalasemia Mayor Di Blud Rsu Cut Meutia Aceh Utara Tahun 2018. AVERROUS: Jurnal Kedokteran dan Kesehatan Malikussaleh. 2018;4(2):62.
18. Origa R, Cinus M, Pilia MP, Gianesin B, Zappu A, Orecchia V, et al. Safety and Efficacy of the New Combination Iron Chelation Regimens in Patients with Transfusion-Dependent Thalassemia and Severe Iron Overload. J Clin Med. 2022;11(7).
19. Entezari S, Haghi SM, Norouzkhani N, Sahebnazar B, Vosoughian F, Akbarzadeh D, et al. Iron Chelators in Treatment of Iron Overload. J Toxicol. 2022;2022.
20. Karami H, Kosaryan M, Hadian Amree A, Darvishi-Khezri H, Mousavi M. Combination Iron Chelation Therapy with Deferiprone and Deferasirox in Iron-Overloaded Patients with Transfusiondependent β-thalassemia Major. Clin Pract. 2017;7(1):912.
21. Kontoghiorghes GJ. The Vital Role Played by Deferiprone in the Transition of Thalassaemia from a Fatal to a Chronic Disease and Challenges in Its Repurposing for Use in Non-Iron-Loaded Diseases. Pharmaceuticals. 2023;16(7).
22. Moukalled NM, Bou-Fakhredin R, Taher AT. Deferasirox: Over a decade of experience in thalassemia. Mediterr J Hematol Infect Dis. 2018;10(1):1–13.
23. Uppal V, Naseem S, Bihana I, Sachdeva MUS, Varma N. Reticulocyte count and its parameters: comparison of automated analyzers, flow cytometry, and manual method. J Hematop. 2020;13(2):89–96.

Published by Universitas Diponegoro.
doi: http://dx.doi.org/.....

Turnitin Artikel

ORIGINALITY REPORT

13%	10%	6%	2%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	ashpublications.org Internet Source	1%
2	repo.stikesicme-jbg.ac.id Internet Source	1%
3	Submitted to University of Western Sydney Student Paper	1%
4	jurnal.globalhealthsciencegroup.com Internet Source	1%
5	www.researchsquare.com Internet Source	1%
6	repository.ju.edu.et Internet Source	1%
7	Husnan Mujiburrahman, Qodri Santosa, Joko Setyono, Nendyah Roestijawati, Lantip Rujito. "Investigating the level of Hba1c and insulin level in β-thalassemia patients", MEDISAINS, 2023 Publication	1%
8	Idris Zubairu Sadiq, Fatima Sadiq Abubakar, Hauwa Salisu Usman, Aliyu Dantani Abdullahi et al. "Thalassemia: Pathophysiology, Diagnosis, and Advances in Treatment", Thalassemia Reports, 2024 Publication	<1%
9	discovery.researcher.life Internet Source	<1%
10	revistas.unisimon.edu.co Internet Source	<1%

11	Submitted to Universitas Airlangga-1 Student Paper	<1%
12	Submitted to Universiti Kebangsaan Malaysia Student Paper	<1%
13	pubmed.ncbi.nlm.nih.gov Internet Source	<1%
14	sciforum.net Internet Source	<1%
15	Antonella Meloni, Laura Pistoia, Cristina Vassalle, Anna Spasiano et al. "Low Vitamin D Levels Are Associated with Increased Cardiac Iron Uptake in Beta-Thalassemia Major", Diagnostics, 2023 Publication	<1%
16	najfnr.com Internet Source	<1%
17	assets.researchsquare.com Internet Source	<1%
18	mdpi-res.com Internet Source	<1%
19	www.scilit.net Internet Source	<1%
20	Lantip Rujito, Tirta Wardana, Joko Mulyanto, Ita Margaretha Nainggolan, Teguh Haryo Sasongko. "Profiling circulating microRNA and regulatory pathways in transfusion-dependent thalassemia and thalassemia trait compared to healthy controls: a preliminary study", ExRNA, 2024 Publication	<1%
21	ijpho.ssu.ac.ir Internet Source	<1%

22	"Abstracts", British Journal of Haematology, 2017 Publication	<1%
23	Lutfia Nurhasanah, Ayu Nissa Ainni, Tri Cahyani Widiastuti, Muh. Husnul Khuluq. "KAJIAN PENGGUNAN OBAT KELASI BESI PADA PASIEN TALASEMIA MAYOR DI RSUD X KEBUMEN TAHUN 2023", Usadha Journal of Pharmacy, 2025 Publication	<1%
24	fdocument.org Internet Source	<1%
25	www.stf.iu.edu.kh Internet Source	<1%
26	ejurnal.ung.ac.id Internet Source	<1%
27	Fisher, Sheila A, Susan J Brunskill, Carolyn Doree, Sarah Gooding, Onima Chowdhury, David J Roberts, and David J Roberts. "Desferrioxamine mesylate for managing transfusional iron overload in people with transfusion-dependent thalassaemia", Cochrane Database of Systematic Reviews Reviews, 2013. Publication	<1%

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	On		

Turnitin Artikel

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6