

## Lampiran 10. Lembar Hasil Analisis Data

**1. Distribusi Karakteristik Nelayan di TPI Puger**

		<b>Usia</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	18-30 tahun	5	5.2	5.2	5.2
	>30 tahun	92	94.8	94.8	100.0
	Total	97	100.0	100.0	

		<b>Jenis Kelamin</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	59	60.8	60.8	60.8
	Perempuan	38	39.2	39.2	100.0
	Total	97	100.0	100.0	

		<b>Pendidikan</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak sekolah	13	13.4	13.4	13.4
	Tidak lulus SD/Sederajat	14	14.4	14.4	27.8
	Lulus SD/Sederajat	32	33.0	33.0	60.8
	Lulus SMP/Sederajat	27	27.8	27.8	88.7
	Lulus SMA/Sederajat	11	11.3	11.3	100.0
	Total	97	100.0	100.0	

		<b>Pendapatan</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	< Rp.2.665.392,00	84	86.6	86.6	86.6
	>= Rp.2.665.392,00	13	13.4	13.4	100.0
	Total	97	100.0	100.0	

**2. Distribusi Faktor Risiko Nelayan di TPI Puger**

		<b>Personal hygiene</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak memadai	66	68.0	68.0	68.0
	Memadai	31	32.0	32.0	100.0
	Total	97	100.0	100.0	

		<b>Masa Kerja</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	>= 5 tahun	82	84.5	84.5	84.5
	< 5 tahun	15	15.5	15.5	100.0
	Total	97	100.0	100.0	

**Obesitas**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Obesitas	53	54.6	54.6	54.6
	Tidak obesitas	44	45.4	45.4	100.0
	Total	97	100.0	100.0	

**1. Distribusi Kejadian Mikosis Superfisialis, Jenisnya, dan Lokasinya****Mikosis Superfisialis**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	10	10.3	10.3	10.3
	Bukan	87	89.7	89.7	100.0
	Total	97	100.0	100.0	

**Jenis Mikosis**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Dermatofitosis	8	8.2	80.0	80.0
	Nondermatofitosis	2	2.1	20.0	100.0
	Total	10	10.3	100.0	
Missing	System	87	89.7		
Total		97	100.0		

**Lokasi Mikosis**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kepala	1	1.0	10.0	10.0
	Badan	8	8.2	80.0	90.0
	Selangkangan	1	1.0	10.0	100.0
	Total	10	10.3	100.0	
Missing	System	87	89.7		
Total		97	100.0		

**2. Hubungan *Personal Hygiene* dengan Kejadian Mikosis Superfisialis*****Personal hygiene* \* Mikosis Superfisialis**

		Mikosis Superfisialis		Total	
		Ya	Bukan		
<i>Personal hygiene</i>	Tidak memadai	Count	8	58	66
		Expected Count	6.8	59.2	66.0
		% within <i>Personal hygiene</i>	12.1%	87.9%	100.0%
	Memadai	Count	2	29	31
		Expected Count	3.2	27.8	31.0
		% within <i>Personal hygiene</i>	6.5%	93.5%	100.0%
Total	Count	10	87	97	
	Expected Count	10.0	87.0	97.0	
	% within <i>Personal hygiene</i>	10.3%	89.7%	100.0%	

Chi-Square Tests					
	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.733 <sup>a</sup>	1	.392		
Continuity Correction <sup>b</sup>	.248	1	.618		
Likelihood Ratio	.791	1	.374		
Fisher's Exact Test				.494	.320
Linear-by-Linear Association	.726	1	.394		
N of Valid Cases	97				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.20.

b. Computed only for a 2x2 table

Risk Estimate			
Value	95% Confidence Interval		Upper
	Lower	Upper	
Odds Ratio for Personal hygiene (Tidak memadai / Memadai)	2.000	.399	10.029
For cohort Mikosis Superfisialis = Ya	1.879	.424	8.333
For cohort Mikosis Superfisialis = Bukan	.939	.826	1.068
N of Valid Cases	97		

### 3. Hubungan Masa Kerja dengan Kejadian Mikosis Superfisialis

#### Masa Kerja \* Mikosis Superfisialis

		Mikosis Superfisialis		Total
		Ya	Bukan	
Masa Kerja >= 5 tahun	Count	10	72	82
	Expected Count	8.5	73.5	82.0
	% within Masa Kerja	12.2%	87.8%	100.0%
< 5 tahun	Count	0	15	15
	Expected Count	1.5	13.5	15.0
	% within Masa Kerja	0.0%	100.0%	100.0%
Total	Count	10	87	97
	Expected Count	10.0	87.0	97.0
	% within Masa Kerja	10.3%	89.7%	100.0%

Chi-Square Tests					
	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.040 <sup>a</sup>	1	.153		
Continuity Correction <sup>b</sup>	.934	1	.334		
Likelihood Ratio	3.564	1	.059		
Fisher's Exact Test				.353	.170
Linear-by-Linear Association	2.019	1	.155		
N of Valid Cases	97				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 1.13.

b. Computed only for a 2x2 table

	Risk Estimate		
	Value	Lower	Upper
For cohort Mikosis Superfisialis = Bukan	.878	.810	.952
N of Valid Cases	97		

#### 4. Hubungan Obesitas dengan Kejadian Mikosis Superfisialis

##### Obesitas \* Mikosis Superfisialis

Obesitas	Obesitas	Count	Mikosis Superfisialis		Total
			Ya	Bukan	
Obesitas	Obesitas	Count	3	50	53
		Expected Count	5.5	47.5	53.0
		% within Obesitas	5.7%	94.3%	100.0%
Tidak obesitas	Tidak obesitas	Count	7	37	44
		Expected Count	4.5	39.5	44.0
		% within Obesitas	15.9%	84.1%	100.0%
Total		Count	10	87	97
		Expected Count	10.0	87.0	97.0
		% within Obesitas	10.3%	89.7%	100.0%

	Chi-Square Tests				
	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.731 <sup>a</sup>	1	.098		
Continuity Correction <sup>b</sup>	1.735	1	.188		
Likelihood Ratio	2.759	1	.097		
Fisher's Exact Test				.178	.094
Linear-by-Linear Association	2.703	1	.100		
N of Valid Cases	97				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.54.

b. Computed only for a 2x2 table

	Risk Estimate		
	Value	Lower	Upper
Odds Ratio for Obesitas (Obesitas / Tidak obesitas)	.317	.077	1.309
For cohort Mikosis Superfisialis = Ya	.356	.098	1.295
For cohort Mikosis Superfisialis = Bukan	1.122	.971	1.296
N of Valid Cases	97		