

The Toxic Metal Arsenic Contamination of the Coastal Aquifers in the North Coast of Java, Indonesia

Chrisna Adhi Suryono

Marine Science Department, Diponegoro University
Jl. Prof. Soedarto, SH. Kampus UNDIP Tembalang, Semarang 50275
Email : chrisna_as@yahoo.com

Abstrak

Pesatnya perkembangan industri dan aktivitas domestic di daerah pantai utara Jawa mendorong untuk perlunya dilakukan penelitian tentang kemungkinan adanya pencemaran logam Arsen (As) di akifer dangkal kota-kota Jakarta, Semarang dan Surabaya. Sebanyak 30 contoh air dari sumur-sumur yang berasal dari zona pemukiman di daerah pesisir dianalisis kandungan logam As dengan menggunakan teknik *Atomic Absorption Spectrophotometer* (AAS). Hasil analisis menunjukkan adanya kandungan logam As sebesar $59.65 \mu\text{g L}^{-1}$ di Jakarta. Rerata dan simpangan baku dari logam As di Jakarta, Semarang and Surabaya adalah 15.47 ± 18.79 , 1.25 ± 2.05 and $0.59 \pm 0.26 \mu\text{g L}^{-1}$. Terdapat korelasi yang nyata antara logam As dan Fe di airtanah. Dari hasil ini dapat disimpulkan bahwa tingkat kandungan logam As dan Fe di daerah akifer pantai utara Jawa masih berada di bawah ambang yang dipersyaratkan oleh *Indonesian Drinking & Domestic Water Quality Standard for Ground Water and WHO's (World Health Organization) Guideline Values for Drinking Water*, kecuali untuk stasion 5 dan 6 dari Jakarta.

Kata kunci: Arsen (As), AAS, akifer pantai

Abstract

A study was conducted to assess the Arsenic (As) metal contamination of the shallow aquifers in Jakarta, Semarang, and Surabaya city of Java and its relation to the highly developed industrial and domestic activities in the coastal region. Arsen was assayed in the waters of 30 wells throughout the terrestrial cities, in residential zones using Atomic Absorption Spectrophotometer (AAS) technique. Analysis of the As level revealed that high As contents were found primarily in Jakarta, to a maximum of $59.65 \mu\text{g L}^{-1}$. The mean and standard deviation of As in Jakarta, Semarang and Surabaya were 15.47 ± 18.79 , 1.25 ± 2.05 and $0.59 \pm 0.26 \mu\text{g L}^{-1}$, respectively. The groundwater As concentration shows a correlation with Fe significantly. It was concluded that the levels of As in some investigated 3 Indonesian metropolis cities were below the maximum allowable concentrations of metals recommended by Indonesian Drinking & Domestic Water Quality Standard for Ground Water and WHO's (*World Health Organization*) Guideline Values for Drinking Water, except for station 5 and 6 of Jakarta.

Keywords: Arsenic (As), AAS, coastal aquifer

INTRODUCTION

Arsenic (As) is one of the most important toxicant that are widely distributed in nature and occur in the form of inorganic and organic compounds. Exposure to inorganic compounds may occur in a variety of ways through certain

industrial effluents, chemical alloys, pesticides, wood preservative agents, combustion of fossil fuels, catalysts, glass, fire retardant, occupational hazards in mining that are anthropogenic and natural sources (Smedly & Kinniburgh, 2002). Arsenic is known to cause arsenicosis due to its manifestation in drinking water, the

most common species being arsenate As (V) and arsenite As (III) (WHO, 2007). Arsenic contamination of groundwater has led to a massive epidemic of arsenic poisoning in Bangladesh (Anawar *et al.*, 2003 and Chowdhury, 2004) and neighbouring countries (Berg *et al.*, 2001 and Nickson *et al.*, 2005). It is estimated that approximately 57 million people in Bangladesh (Yu, *et al.*, 2003) and 11 million people in Vietnam (Berg *et al.*, 2001) are drinking groundwater with arsenic concentrations elevated above the World Health Organization's standard of 10 parts per billion (WHO, 1993). This tragedy was occurring as the high arsenic levels present in drinking water. Bangladesh maintains an allowable limit of 50 ppb, and the limit in Canada is 25 ppb (Nordstrom, 2002). The arsenic found in Bengale and Canada drinking water is present naturally in the sediment but is released only by humans, primarily from the digging of shallow wells. Similarly unacceptably high arsenic levels are found in shallow-well domestic drinking water in the American Midwest (Kim, *et al.*, 2002).

Recently there has been increasing anxieties concerning arsenic related problems. Occurrence of arsenic contamination has been reported worldwide, for example, Mongolia (Gong *et al.*, 2006), USA (Welch *et al.*, 2000; Kim *et al.*, 2002 and Hudak, 2008;), Bangladesh (Anawar *et al.*, 2003 and Chowdhury, 2004), Taiwan (Chen *et al.*, 1994), Spain (García-Sánchez *et al.*, 2005), *et cetera*. In Indonesia, no significant As contamination has been reported yet. However, local contaminations in soils, sea and groundwaters with high As and Hg levels have been frequently revealed particularly around Buyat Bay mining areas (Khalik and Wulandari, 2004 and KLH, 2004).

Most of the water requirement for metropolis city such as Jakarta, Semarang, Surabaya, is met from surface and groundwater supplies piped in from outside the city limits. However, the case for many cities, a certain proportion of the population, usually the poorest, is forced to rely on dug and shallow wells within the urban area. A study was therefore carried




Figure 1. The map of sampling site locations.

out to gain an idea of the inorganic quality of the water in the wells penetrating the shallow aquifer below the city. This research aim to examine the As content of shallow aquifer in coastal urban areas of Jakarta, Semarang and Surabaya and assess how far they conform with the WHO/Indonesia recommendations for drinking water.

MATERIAL AND METHODS

Water collections

A total of thirty samples were collected at each of the stations. Water samples used for As analyses were collected from the residential area shallow aquifer of the north shore of Jakarta, Semarang, and Surabaya metropolis during July-August, 2007 (Fig.1). Shallow aquifer waters are the only sources for drinking water in the study area, however some people uses these waters directly for domestic and drinking and not through the water supplies system. The water samples were collected in 1000 mL plastic bottles. Then, bottles were properly labeled and tightly sealed. All the water samples were brought to the laboratory for performing water quality tests and heavy metal analyses.

Arsenic metal analysis.

The water samples were acidified with aliquot of HNO₃. About 100 mL each of the wellmixed acidified water was digested with 5 mL HCl. The digested samples were then analyzed for As and Fe using atomic absorption spectrometer Perkin Elmer Model 5000 (Kremling *et al.*, 1998). A statistical analysis was conducted among the standard parameters using Linear correlation method. This was performed to know whether any relationship among the parameters existed or not.

Statistical analysis

The value was used to analyze correlations between As and Fe. ANOVA analysis was made among locations of heavy metal content. All the statistical

methods were done using the software Minitab ver.15. Analysis of variance ANOVA was employed after logarithmic conversion when necessary to detect significant differences among mean.

RESULT AND DISCUSSION

The appearance of arsenic in groundwater and drinking water is a worldwide concern since it has been found that arsenic is one of the most toxic and carcinogenic chemical elements. In the present study As metal concentration was ranging from 0.59 ± 0.26 to 15.47 ± 18.79 μgL^{-1} respectively in three different cities of Java north shore. The maximum As concentration was observed in Jakarta (59.65 μgL^{-1}) and minimum was observed in Semarang and Surabaya (undetected). Statistically, there were significant differences among locations, Jakarta, Semarang and Surabaya on quantity of As content. Both Semarang and Surabaya were no significant different on As (Table 1). This condition could be explained that first, arsenic can enter the water supply from natural deposits in the earth or from industrial and agricultural pollution.

It is a natural element used for a variety of purposes within industry and agriculture. It is also a byproduct of copper smelting, mining, and coal burning. Industries in Jakarta were suspected release much more arsenic into the environment every year than those of Semarang and Surabaya. Once released, arsenic remains in the environment for a long time. However, in the present results of this study when it was compare to the previous results in Buyat Bay for Drinking Water (KLH, 2004), it was lower. Because it may be stated that the environmental parameters like salinity, temperature, dissolved oxygen and pH also have some effect on the accumulation of trace metals.

Among the parameters the salinity played major role in the metal accumulation (Rajan, 1987). Figure 2 showed a positive and significant correlation between As and Fe (Chrisna *et al.*, 2007 *in press*) in coastal aquifer of Java

Table 1. Range, mean and standard deviation of As content (g L⁻¹) in coastal aquifer of Java north shore.

Sampling Station	As concentration (µg L ⁻¹) in:		
	Jakarta	Semarang	Surabaya
ST1	7.5	0.54	0.25
ST2	33.65	0.13	0.39
ST3	5.76	<0.001	0.40
ST4	7.58	1.63	1.11
ST5	21.07	0.50	0.66
ST6	59.65	6.21	0.89
ST7	1.90	3.42	0.43
ST8	<0.001	0.03	0.51
ST9	16.9	<0.001	0.54
ST10	0.69	<0.001	0.76
Range	0 – 59.65	0 – 6.21	0.25 – 1.11
Mean and Standard deviation	15.47 ± 18.79 a	1.25 ± 2.05 b	0.59 ± 0.26 b

Note: The difference of alphabet showed significant difference




Figure 2. Relation of Arsenic and iron in coastal aquifer (p=0.005)

north shore. As concentrations correlate well with concentration of dissolved Fe. It means that the higher the dissolved Fe, the arsenic concentration increase. Many authors have reported the correlations between As and Fe in a variety of environments, often with the explanation that As adsorbs onto Fe (hydroxides) under oxidizing conditions and is then released when the (hydroxide) undergoes reductive dissolution (Zheng *et al.*, 2004; Mirlean *et al.*, 2006;). Bhattacharya *et al.* (1997) reported that arsenic is released when arsenic-rich

iron oxyhydroxides are produced in anoxic groundwater. In addition, Nickson *et al.* (1998) stated that arsenic derived from the oxidation of arsenic-rich pyrite in the aquifer sediments as atmospheric oxygen invades the aquifer in response to a lowering of the water level by abstraction. Previous investigations in Bangladesh have also suggested that arsenic may enter the aquifer associated with iron oxide coatings on the sediment grains and when conditions become reduced, the arsenic is released to the groundwater together with

Fe²⁺ (McArthur *et al.*, 2001; Smedly and Kinniburgh, 2002). Bone *et al.* (2006) reported that strong correlation between As and Fe provides solid evidence that As is linked to the Fe redox cycle.

CONCLUSION

This experiment has shown that the entire water sample meets the limits of WHO and Indonesian Drinking & Domestic Water Quality Standard for Ground Water, except for station 5. This study confirmed a significant relationship between the arsenic concentration and iron concentration in coastal aquifer of Java north shore. It means that coastal aquifer of Java north shore is not polluted. With regard to mitigation, shallow aquifer with low concentrations of arsenic in water which are continuing to be used need to be monitored over time by authorities concerned to ensure that concentrations of arsenic in water of these shallow aquifer remain at an acceptable level over time.

REFERENCES

- Anawar, H.M., J. Akai, K. Komaki, H. Terao, T. Yoshioka, T. Ishizuka, S. Safiullah, and K. Kato. 2003. Geochemical occurrence of arsenic in groundwater of Bangladesh: Sources and mobilization processes. *J. Geochem. Explor.* 77:109–131.
- Anonymous. 1997. Water quality. Ohio Department of Natural Resources. Division of Water. 3 p.
- Anonymous. 2003. Arsenic and Drinking Water from Private Wells. NSF International.
- Anonymous. 2008. Key to Metals AG. All Rights Reserved.
- Bautista, M.A. and A.K. Pradhan. 1995. Iron and Nickel Abundances in H-II Regions and Supernova Remnants.
- Berg, M., Tran, H.C., Nguyen, T.C., Pham, H.V., Schertenleib, R., Giger, W., 2001. Arsenic contamination of groundwater and drinking water in Vietnam: a human health threat. *Environ. Sci. Technol.* 35, 2621–2626.
- Bhattacharya, P., D. Chatterjee, and G. Jacks. 1997. Occurrence of arsenic-contaminated groundwater in alluvial aquifers from delta plains, eastern India: Options for safe drinking water supply. *Water Resour. Dev.* 13:79–92.
- Bone, E.S., M.E. Gonnee, and M.A. Charette. 2006. Cycling of Arsenic in Coastal Aquifer. *Environ. Sci. Technol.* 40:3273–3278
- Chen, S.L., S.R. Dzung, M.H. Yang, K.H. Chiu, G.M. Shieh, and C.M. Wai. 1994. Arsenic species in groundwaters of the blackfoot disease area, Taiwan. *Environ. Sci. Technol.* 28:877–881.
- Chowdhury, A. M. R. 2004. Arsenic crisis in Bangladesh. *Sci. Am.* 291:86-91.
- Chrisna A. S, Baskoro, R, Agus, S and Tyas, S. B., 2007. Physico-chemical Characteristics and Heavy Metal Contents in Shallow Groundwater of Semarang Coastal Region. (in press). *Ilmu Kelautan* (12) 3:
- García-Sánchez, A., A. Moyano and P. Mayorga. 2005. High arsenic contents in groundwater of central Spain. *Environ. Geol.* Vol. 47(6):847–854
- Gong, Z., X. Lu, C. Watt, B. Wen, B. He, J. Mumford, Z. Ning, Y. Xia and X. C. Le. 2006. Speciation analysis of arsenic in groundwater from Inner Mongolia with an emphasis on acid-leachable particulate arsenic. *Analytica Chimica Acta* Vol. 555 (1):181-187
- Hudak, P.F. 2008. Distribution of arsenic concentrations in groundwater of the Seymour Aquifer, Texas, USA. *Int. J. of Environ. Health Research*, Vol. 18 (1): 79 – 82
- Khalik, A. and F. Wulandari. 2004. Buyat Bay tests show high mercury levels. *Jakarta Post* July 30, 2004
- Kim, M. J., J. Nriagu, and S. Haack. 2002. Arsenic species and chemistry in groundwater of southeast Michigan. *Environ. Pollut.* 120:379-390.
- KLH .2004. REPORT. Environmental Quality Assessment of Buyat Bay and Totok Bay. Kementerian Lingkungan Hidup. 41 p
- Kremling, K., Andreae, M.O., Brugmann, L., van den Berg, C, M, G., Prange, A., Schirmacher., Koroleff, F., and Kuss, J., 1998. Determination of Trace Elements in Method of Seawater Analysis. Wiley-Vch. Toronto. 254 – 364 pp

- McArthur, J.M., Ravenscroft, P., Safiulla, S., Thirlwall, M.F., 2001. Arsenic in groundwater: testing pollution mechanisms for sedimentary aquifers in Bangladesh. *Water Resour. Res.* 37, 109–117.
- Mirlean, N.; V. E. Andrus, P. Baisch, G.Griep, and M.R.Casartelli. 2006. Arsenic pollution in Patos Lagoon estuarine sediments, Brazil. *Mar. Pollut. Bull.* Vol. 46: 1480-1484.
- Nickson, R., J. McArthur, W. Burgess, K.M. Ahmed, P. Ravenscroft and M. Rahman. 1998. Arsenic poisoning of Bangladesh groundwater. *Nature* Vol.395: 338
- Nickson, R.T., J.M. McArthur, B. Shrestha, T.O. Kyaw-Myint, and D. Lowry. 2005. Arsenic and other drinking water quality issues, Muzaffargarh District, Pakistan. *Applied Geochemistry* 20: 55–68
- Nordstrom, D.K. 2002. Worldwide Occurrences of Arsenic in Ground Water. *Science* Vol.296: 2143-2144
- Rajan, A , 1987. Studes on *Donax cuneatus* (Linnaeus) (Mollusca: Bivalvia: Donacidae) from Polto Novo waters. PhD Thesis, Annamalai University. In Shanmugam, A., C. Palpandi and K. Kesavan. 2007. Bioaccumulation of Some Trace Metals (Mg, Fe, Zn, Cu) from Begger's Bowl *Cymbium melo* (Solander, 1786) (A Marine Neogastropod). *Research J. of Environ. Sci.* 1 (4): 191-195
- Smedley, P.L. and DG Kinniburgh, 2002. Areview of the source, behaviour and distribution of arsenic in natural waters. *Applied Geochem.*, 17: 51 7-568.
- Welch, A.H., D.B.Westjohn, D.R. Helsel and R.B. Wanty. 2000. Arsenic in ground water of the United States: occurrence and geochemistry. *Ground Water*. Vol. 38 (4):589-604.
- WHO, 1993. Guidelines for Drinking-water Quality, 2nd ed., vol. 1. Recommendations, Geneva.
- WHO, 2007. Arsenic in Drinking Water .
- Yu, W.H., Harvey, C.M., Harvey, C.F., 2003. Arsenic in groundwater in Bangladesh: a
- Zheng, Y.; M. Stute, A. van Geen, I. Gavrieli, R Dhar, H.J. Simpson, P. Schlosser, M.K.Ahmed. 2004. Redox control of Arsenic mobilization in Bangladesh groundwater. *Appl. Geochem.* Vol.19: 201-214.