

manuscrip turnitin.docx

by sitinurjannnah@gmail.com sitinurjannnah@gmail.com

Submission date: 04-Mar-2026 05:15PM (UTC+0700)

Submission ID: 2894120797

File name: manuscrip_turnitin.docx (124.55K)

Word count: 4004

Character count: 26036

Is Insurance Membership Type Associated with Disease Claims under Non-Capitation Payment at Primary Healthcare Facilities? (Analysis of BPJS Kesehatan Sample Data, 2015–2023)

Mohammad Yusuf Setiawan pendanaan, Achmad Djunawan*, Enny Mar'atus Sholihah

Abstract

Introduction: Primary healthcare plays a central role in Indonesia's National Health Insurance system in managing population health needs. Understanding morbidity patterns and their relationship with insurance membership categories is essential for service planning and financial sustainability. This study aimed to analyze disease claim patterns at non-capitation primary healthcare facilities and examine their association with insurance membership types between 2015 and 2023.

Methods: A retrospective cross-sectional study was conducted using secondary administrative claims data from primary healthcare facilities under the National Health Insurance scheme. The dataset consisted of 201,297 non-capitation service records collected between 2015 and 2023. Variables included diagnosis codes, insurance membership type, sex, and facility characteristics. Data were analyzed descriptively to identify morbidity patterns and inferentially using chi-square tests to assess associations between membership categories and disease claims.

Results: Non-communicable diseases, particularly type 2 diabetes mellitus and essential hypertension, were the most frequently reported diagnoses across all membership categories. Maternal and reproductive health conditions also ranked among the top diagnoses, especially among government-subsidized participants. A statistically significant association was found between insurance membership type and morbidity pattern ($p < 0.001$), although the strength of the association was weak.

Conclusion: Morbidity patterns in primary healthcare reflect a dual burden of chronic and maternal conditions and vary across insurance membership categories. These findings highlight the need for membership-based intervention strategies and strengthened chronic disease management to ensure sustainability within the National Health Insurance system.

Keywords: chronic disease, health insurance, morbidity pattern, primary healthcare, universal health coverage

Introduction

Primary Health Care Facilities (PHCFs) serve as the frontline of health service delivery within Indonesia's National Health Insurance system (Jaminan Kesehatan Nasional, JKN). Ideally, PHCFs are expected to deliver efficient and equitable services, particularly for non-specialistic and chronic conditions, to minimize unnecessary referrals to higher-level health facilities. Financing at the primary care level is conducted through capitation and non-capitation payment mechanisms, which should be aligned with participant characteristics and morbidity burden rather than merely enrollment numbers. A well-designed financing system can encourage early detection, comprehensive disease management, and continuous monitoring of population health outcomes. Consequently, morbidity patterns observed at PHCFs may reflect the overall health status of the insured population in a given area.

Despite the expansion of JKN coverage, significant challenges remain in managing morbidity and financing at the primary care level. National data indicate that non-capitation expenditure at PHCFs reached approximately IDR 1.6 trillion in 2022, while non-INA-CBG expenditure at referral facilities amounted to IDR 2.3 trillion, highlighting inefficiencies in primary care management (Nushratu, 2023). By September 2024, JKN coverage had reached 98.19% of the population, equivalent to approximately 277 million individuals (Alatas, 2024).

Furthermore, national morbidity data published in the *Indonesian Population Morbidity Table* demonstrate a substantial disease burden across JKN participants (Alatas, 2024). These trends suggest that morbidity at the primary care level has not been optimally addressed through existing financing arrangements.

Several previous studies have examined morbidity or financing aspects in PHCFs, although most focused on specific diseases or regions. A study conducted in West Java reported that among 9,056 BPJS participants receiving non-capitation services for hypertension at PHCFs during 2015–2016, the majority were female and belonged to the self-employed membership segment (PBPU) with Class III premiums (Dewi et al., 2021). Another study in Central Java found that the morbidity rate in 2018 reached 15.15%, exceeding the national average of 13.91% (Rosanti & Budiantara, 2020). Research on pulmonary tuberculosis patients also revealed that JKN membership type was significantly associated with the utilization of PHCF services (Rahman & Hidayat, 2024). However, these studies rarely integrated morbidity patterns with non-capitation financing mechanisms and membership segmentation.

This situation reveals a clear research gap in the existing literature. To date, few quantitative studies have systematically analyzed disease claim or morbidity patterns among PHCF patients receiving non-capitation financing based on different JKN membership categories. Most previous research has emphasized capitation-based payments or general service utilization, while the role of non-capitation financing in shaping morbidity patterns remains underexplored. Additionally, many studies focus on specific diseases, such as hypertension, rather than examining overall morbidity distributions. The limited integration of morbidity, non-capitation financing, and membership type restricts the development of evidence-based policies for primary care financing.

Understanding morbidity patterns within non-capitation financing is particularly important given the heterogeneity of JKN membership categories. Government-subsidized members (PBI), salaried workers (PPU), and self-employed participants (PBPU) differ in socioeconomic characteristics, access to care, and health risks, which may result in distinct morbidity and claim patterns. Ineffective alignment between financing mechanisms and morbidity burden may lead to inefficient resource allocation, excessive referrals, and inequities in service delivery. These challenges have been linked to broader financial sustainability concerns within the JKN system managed by BPJS Kesehatan, particularly amid the increasing dominance of chronic diseases (Jatmiko, 2018).

Therefore, this study aims to analyze disease claim patterns among patients treated at PHCFs under non-capitation financing based on JKN membership categories. Using a quantitative approach, this study examines the distribution of disease diagnoses, compares claim patterns across membership groups, and assesses the relationship between membership type and non-capitation disease claims at the primary care level. By integrating morbidity patterns, financing mechanisms, and membership segmentation, this study seeks to address limitations in previous research. The findings are expected to contribute to the literature on primary health care financing and support evidence-based policy formulation to strengthen primary care services and advance high-quality Universal Health Coverage in Indonesia.

Methods

Study Design and Setting

This study employed an analytic quantitative approach using secondary data obtained from the Indonesian National Health Insurance system. A cross-sectional design was applied, as all variables were analyzed at a single point in time to examine differences and associations between variables. The study utilized national-level data from Primary Health Care Facilities (PHCFs) across Indonesia. Secondary data were sourced from the central database of BPJS Kesehatan, covering services delivered at PHCFs under non-capitation financing schemes.

Study Period

The study used secondary data spanning from 2015 to 2024, encompassing nearly a decade of PHCF service records. The research implementation, including data acquisition, statistical analysis, and reporting, was scheduled between October and December 2025. This period allowed sufficient time for data cleaning, classification, and comprehensive analysis. The extended timeframe of the dataset enabled a robust assessment of morbidity patterns across membership categories.

Population, Sampling Method, and Unit of Analysis

The study population consisted of all BPJS Kesehatan participants who received PHCF services financed through non-capitation mechanisms during the study period. A total sampling method was applied, as all eligible records meeting the inclusion criteria were included in the analysis. The final dataset comprised 201,297 individuals, each representing one unit of analysis. Each individual record included information on primary diagnosis and membership category, allowing for comprehensive population-level analysis without sample size estimation or power calculation.

Variables and Operational Definitions

The independent variable in this study was BPJS Kesehatan membership type, categorized as government-subsidized members (PBI—central and local government), salaried workers (PPU), self-employed members (PBPU), and non-workers. The dependent variable was disease claim patterns at PHCFs under non-capitation financing, representing morbidity. Disease claims were identified using the International Classification of Diseases, 10th Revision (ICD-10). Morbidity was measured as the frequency and proportion of ICD-10 diagnosis codes across membership categories, using nominal scales for all variables.

Data Collection Procedures and Research Instruments

Data collection began with a formal request for access to secondary data submitted to BPJS Kesehatan. Upon approval, datasets were received in digital format (.dta files) containing information on diagnoses (ICD-10 codes), membership type, financing mechanism, age, and geographic region. Diagnoses were subsequently classified according to ICD-10 Chapters I–XXII to facilitate standardized morbidity grouping. The primary research instrument was a structured secondary data extraction form, supported by the official ICD-10 classification published by the World Health Organization.

Data Processing and Statistical Analysis

Data processing and analysis were conducted using Google Colaboratory as the primary analytical platform. Descriptive statistical analysis was performed to summarize the distribution of diagnoses and membership categories using frequencies and percentages. Inferential analysis was conducted using the chi-square test to assess differences in morbidity patterns across BPJS Kesehatan membership types. To examine the strength of association between membership type and disease claims, contingency coefficient analysis was applied. All results were presented in tabular and graphical formats, with analysis focused on identifying morbidity pattern variations among membership segments.

Ethical Considerations

This study involved the use of secondary administrative health data obtained from the Indonesian National Health Insurance system. Ethical approval for this research was granted by the Health Research Ethics Committee of STIKES Yayasan Rumah Sakit Dr. Soetomo (Reference No. **KEPK/YRSDS/075/XII/2025**). The ethics committee reviewed and approved the research protocol in accordance with the principles of human rights protection, seven World Health Organization (WHO) standards, and the 2016 Council for International Organizations of Medical Sciences (CIOMS) guidelines.

All data used in this study were anonymized and did not contain personally identifiable information. Therefore, the requirement for individual informed consent was waived by the

ethics committee. All research procedures were conducted in compliance with national regulations and international ethical standards for research involving human data.

Results

General Characteristics of the Data

The initial stage of analysis involved an overview of the dataset to obtain a comprehensive understanding of its structure and characteristics. This study used secondary data on patient services at Primary Health Care Facilities (PHCFs) obtained from BPJS Kesehatan, covering the period from 2015 to 2023. The dataset, referred to as non-capitation data, consisted of 201,297 observations and was organized in a tabular format containing participant characteristics, diagnosis information, and service-related variables. Additional membership data were merged using PSTV01 as the primary key to obtain variables such as sex, marital status, and membership class. This process ensured data consistency and completeness prior to statistical analysis.

Sex Distribution of Patients

Figure 8 presents the distribution of patients by sex among PHCF users receiving non-capitation services. The majority of patients were female, accounting for 147,846 individuals (73.4%), while male patients numbered 53,451 (26.6%). This indicates a clear predominance of female patients utilizing primary care services under non-capitation financing. The observed distribution suggests that women accessed PHCF services more frequently than men during the study period. This imbalance was considered in interpreting subsequent morbidity patterns.

Marital Status of Patients

The distribution of marital status is illustrated in Figure 9. Most patients were married, totaling 159,664 individuals (79.3%), followed by unmarried patients at 22,152 (11.0%) and divorced patients at 15,899 (7.9%). A small proportion of records (1.8%) had undefined marital status. This distribution indicates that married individuals constituted the largest proportion of PHCF service users. Marital status was therefore considered a relevant demographic characteristic in describing the patient population.

Membership Class Distribution

Figure 10 shows the distribution of BPJS Kesehatan membership classes among patients receiving non-capitation services. More than half of the patients were enrolled in class III, accounting for 104,591 individuals (52.0%). Patients in class I and class II represented 26.9% and 21.1% of the sample, respectively. Missing data for membership class were minimal (0.1%). These findings indicate a predominance of lower-class membership among PHCF users under non-capitation schemes.

Ownership of Primary Health Care Facilities

Figure 11 displays the distribution of PHCF ownership utilized by patients. The majority of patients accessed private PHCFs, accounting for 140,744 cases (69.9%). Government-owned facilities at the district or municipal level contributed 26.1% of cases, while facilities owned by provincial governments and state-owned enterprises accounted for much smaller proportions. Facilities operated by the military and police contributed only a negligible number of cases. This distribution highlights the dominant role of private and local government PHCFs in delivering non-capitation services.

Geographic Distribution by City and Province

The distribution of PHCF locations by city is presented in Figure 12. The highest utilization was observed in Semarang City (20.6%), followed by Malang City (11.7%). Several other districts and cities showed relatively balanced proportions ranging between 9% and 10%. Figure 13 presents the distribution by province, showing that Central Java dominated with

32.6%, followed by East Java (24.3%) and West Java (13.8%). These findings indicate a strong concentration of data from provinces located on Java Island.

Disease Claims Profile


Disease claims in this study represent diagnoses recorded during PHCF visits financed through non-capitation mechanisms. Figure 14 illustrates the ten most frequent ICD-10 diagnoses among patients. Type 2 diabetes mellitus (E11) was the most common diagnosis, accounting for 66,861 cases (37.8%), followed by essential hypertension (I10) with 49,013 cases (27.7%). Several maternal and reproductive health-related diagnoses, such as supervision of normal pregnancy (Z34) and contraceptive management (Z30), were also observed, although at lower proportions. Overall, the distribution indicates a predominance of non-communicable chronic diseases.

Distribution of BPJS Kesehatan Membership Types

Figure 15 shows the distribution of BPJS Kesehatan membership types among patients. The largest proportion consisted of salaried workers (PPU), totaling 55,573 participants (27.6%). This was followed by government-subsidized members funded by the national budget (PBI APBN) at 24.5% and self-employed members (PBPU) at 23.6%. Non-workers accounted for 15.1%, while PBI APBD represented the smallest group at 9.3%. This distribution reflects a dominance of formal workers and independent contributors among PHCF users.

Morbidity Patterns by Membership Type

Figure 16 presents the ten most common diseases stratified by BPJS Kesehatan membership type. Type 2 diabetes mellitus (E11) and essential hypertension (I10) were the most frequent diagnoses across all membership categories, particularly among PPU and PBPU participants. Maternal health-related conditions, such as supervision of normal pregnancy (Z34) and contraceptive management (Z30), were more prominent among PBI APBN members. Other chronic conditions, including unspecified diabetes mellitus (E14) and type 1 diabetes mellitus (E10), showed relatively consistent distributions across membership types. These patterns indicate variation in morbidity profiles according to insurance membership.



Association Between Membership Type and Morbidity

Morbidity	Jenis Kepesertaan					Total
	BUKAN PEKERJA	PBI APBN	PBI APBD	PBPU	PPU	

	n	%	n	%	n	%	n	%	n	%	n	%
E11 Type 2 diabetes mellitus	15.273	22,84	11.098	16,6	3.635	5,44	19.765	29,56	17.090	25,56	66.861	100
I10 Essential (primary) hypertension	10.461	21,34	11.181	22,81	3.304	6,74	12.081	24,65	11.986	24,45	49.013	100
Z34 Supervision of normal pregnancy	20	0,18	3.643	32,31	1.755	15,56	1.885	16,72	3.973	35,23	11.276	100
Z30 Contraceptive management	11	0,10	4.689	43,52	1.579	14,66	1.149	10,66	3.346	31,06	10.774	100
E14 Unspecified diabetes mellitus	1.903	18,31	1.751	16,85	372	3,58	2.682	25,8	3.686	35,46	10.394	100
Z39 Postpartum care and examination	9	0,13	2.581	36,39	1.297	18,29	1.212	17,09	1.993	28,10	7.092	100
O80 Single spontaneous delivery	8	0,12	2.634	38,88	1.304	19,25	1.181	17,43	1.648	24,32	6.775	100
E10 Type 1 diabetes mellitus	1.334	20,84	1.034	16,15	359	5,61	1.657	25,88	2.018	31,52	6.402	100
Z36 Antenatal screening	4	0,07	1.805	31,63	988	17,31	906	15,88	2.004	35,11	5.707	100
I15 Secondary hypertension	356	12,71	709	25,32	207	7,39	621	22,18	907	32,39	2.800	100
Total	29.379	16,5894	41.125	23,2221	14.800	8,35714	43.139	24,3593	48.651	27,4718	177.094	100

The association between BPJS Kesehatan membership type and disease claims was examined using chi-square analysis. Table 2 presents the cross-tabulation of the ten most frequent diseases by membership category. The chi-square test yielded a statistic of $\chi^2 = 22,541.61$ with 36 degrees of freedom and a p-value < 0.001, indicating a statistically significant association between disease diagnosis and membership type. However, the Cramer's V value of 0.1784 suggests that the strength of this association was weak. This indicates that while morbidity patterns differed significantly across membership types, the magnitude of the relationship was relatively small.

Discussion

The findings of this study indicate that non-capitation claims at primary healthcare facilities (FKTP) in Indonesia between 2015 and 2023 were predominantly driven by non-communicable diseases, particularly type 2 diabetes mellitus (E11) and essential hypertension (I10). This pattern reflects the ongoing epidemiological transition in Indonesia, where disease burden has shifted from infectious diseases to chronic conditions associated with lifestyle changes and urbanization. Similar trends have been reported in national and international studies, which document a rapid increase in diabetes and hypertension prevalence in middle-income countries (Nugraheni et al., 2020). The dominance of chronic disease cases also explains the high volume of repeat visits to FKTP for long-term disease management. These findings reinforce the critical role of primary care in controlling non-communicable diseases within the JKN system.

The higher proportion of female patients observed in this study suggests gender-based differences in healthcare utilization at the primary care level. Previous research consistently shows that women tend to have better health-seeking behavior and higher utilization of preventive and routine healthcare services compared to men. In the Indonesian context, this pattern is further reinforced by the availability of reproductive and maternal health services delivered through FKTP. Studies on primary healthcare utilization in Indonesia have demonstrated that women are more likely to access services related to screening, chronic disease follow-up, and maternal care (Djunawan, If'al Lillah, et al., 2022). Consequently, gender composition should be considered when interpreting morbidity patterns in primary healthcare data.

Maternal and reproductive health diagnoses, such as supervision of normal pregnancy (Z34), contraceptive management (Z30), spontaneous delivery (O80), and postpartum care (Z39),

appeared prominently among the top diagnoses. This finding highlights the strategic role of FKTP as the primary gateway for maternal and child health services under the JKN scheme. Previous evaluations of JKN implementation have shown that health insurance coverage encourages increased utilization of antenatal and delivery services due to reduced financial barriers. Evidence from Indonesia also indicates that the expansion of health insurance coverage has contributed to improved access to skilled birth attendance and facility-based deliveries (Djunawan & Haksama, 2015). The prominence of maternal diagnoses in this study therefore reflects the effectiveness of subsidized insurance in supporting maternal healthcare utilization.

The presence of multiple diabetes-related codes (E10, E11, and E14) illustrates the broad spectrum of metabolic disorders managed at the primary care level. Prior studies have noted that glycemic and blood pressure control in FKTP remains challenging due to issues such as patient adherence and limitations in long-term monitoring. National programs such as Prolanis were designed to strengthen chronic disease management at FKTP, yet their implementation remains uneven across regions. Research on healthcare utilization patterns indicates that individuals with more stable insurance coverage are more likely to engage in routine chronic disease management. This may explain why diabetes and hypertension dominate non-capitation claims within the JKN system.

Regarding insurance membership, the dominance of Pekerja Penerima Upah (PPU), followed by PBI APBN and PBPU, reflects the broad coverage of JKN across both formal workers and subsidized populations. This structure aligns with the goal of achieving universal health coverage by integrating formal, informal, and vulnerable population groups. Previous studies have shown that possession of health insurance significantly increases primary care utilization, particularly in urban settings (Djunawan, 2018). The substantial presence of subsidized participants in FKTP utilization underscores the importance of government financing in expanding access to healthcare for low-income populations. From a policy perspective, these findings highlight the need to align service capacity with membership composition.

The statistically significant association between insurance membership and morbidity patterns indicates that different participant groups utilize FKTP services for different health needs. Chronic diseases were more prevalent among PPU and PBPU participants, while maternal health services were predominantly utilized by PBI APBN and PBI APBD participants. This pattern supports the social determinants of health framework, which emphasizes the role of socioeconomic status and financial protection in shaping disease patterns (Nugraheni et al., 2020). However, the relatively low Cramer's V value suggests that insurance membership alone explains only a small proportion of morbidity variation. Other factors such as age, gender, lifestyle, and geographic location likely play substantial roles.

Several limitations should be considered when interpreting the findings of this study. First, the analysis relied on secondary administrative data, which limits the availability of clinical details and behavioral risk factors. Second, the cross-sectional nature of the data aggregation prevents causal inference regarding the relationship between insurance membership and morbidity patterns. Third, potential misclassification of diagnoses and incomplete recording in routine claims data may have influenced the distribution of disease categories. Despite these limitations, the large sample size and long observation period provide robust insights into utilization patterns within the JKN system.

Overall, the study demonstrates that primary healthcare utilization under JKN is shaped by both epidemiological transition and insurance membership structure. The coexistence of high non-communicable disease burden and substantial maternal service utilization places significant demands on FKTP capacity. Strengthening promotive and preventive services, particularly for chronic disease risk factors, is essential to control future healthcare costs. At the same time, continued support for maternal and reproductive health services remains crucial

for protecting vulnerable populations. These findings underscore the importance of targeted primary care interventions to ensure the sustainability and effectiveness of Indonesia's national health insurance system (Djunawan, 2019).

Conclusion

This study demonstrates that morbidity patterns in non-capitation primary healthcare facilities under Indonesia's National Health Insurance (JKN) system are predominantly driven by non-communicable diseases, particularly type 2 diabetes mellitus and essential hypertension, while maternal and reproductive health services remain a substantial component of care. The coexistence of chronic disease management and maternal health utilization reflects a dual service burden at the primary care level.

Significant differences in morbidity distribution were observed across insurance membership categories. Chronic conditions were more prominent among salaried and self-employed contributors (PPU and PBPU), whereas maternal and reproductive services were concentrated among subsidized participants (PBI). Although the statistical association was significant, its modest strength indicates that insurance membership interacts with broader demographic and socioeconomic determinants in shaping healthcare utilization.

The relevance of this study lies in its empirical demonstration that membership segmentation within a universal health coverage system influences service demand patterns. Using nine years of administrative claims data, the findings provide actionable evidence for capacity planning, risk pooling considerations, and targeted preventive strategies within JKN.

Policy implications include strengthening structured chronic disease management for economically active contributors, sustaining comprehensive maternal services for subsidized populations, and implementing membership-based intervention design. Furthermore, leveraging longitudinal and predictive analytics on JKN data is essential to anticipate future morbidity trends and ensure financial sustainability. These findings contribute to the broader discourse on how insurance design shapes primary healthcare performance in middle-income countries pursuing universal health coverage.

Ethics approval

This study involved the analysis of secondary administrative data derived from Indonesia's National Health Insurance (JKN) primary healthcare claims database. Ethical approval was obtained from the Health Research Ethics Committee of STIKES Yayasan Rumah Sakit Dr. Soetomo (Reference No. KEPK/YRSDS/075/XII/2025). The research protocol was reviewed and approved in accordance with national ethical standards and international guidelines for research involving human data.

The dataset used in this study consisted of anonymized administrative claims records without personally identifiable information. As the study relied exclusively on de-identified secondary data, the requirement for individual informed consent was formally waived by the ethics committee. All procedures were conducted in compliance with applicable ethical regulations governing research involving human participants and health data.

Availability of data and materials

The data that support the findings of this study are available from BPJS Kesehatan (Indonesia's National Health Insurance Agency). However, restrictions apply to the availability of these data because they were accessed under institutional permission and contain administrative health claims information that may involve sensitive individual-level records. Therefore, the datasets are not publicly available.

Data may be made available from the corresponding author upon reasonable request and with permission from BPJS Kesehatan, subject to applicable data protection regulations and institutional policies.

Acknowledgment

Not applicable.

25

Funding

This research was funded independently by the authors. No external funding was received for the design of the study, data collection, analysis, interpretation of data, or writing of the manuscript. The funding source had no role in the study design, data management, analysis, interpretation, or preparation of the manuscript.

Author Contribution

MYS provided funding support for the study. AD designed the study, performed data analysis, drafted the manuscript, and conducted critical revisions of the intellectual content. EMS was responsible for data collection and data preparation. All authors read and approved the final manuscript.

References

ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|---|
| <div style="background-color: red; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">1</div> | <p>Eka Wilda Faida, Diah Wijayanti Sutha, Alfina Aisatus Saadah. "Association Between Dimension of Intelligence and Work Performance among Medical Data Recorders: A Cross-sectional Study", Journal of Public Health Sciences, 2025</p> <p>Publication</p> | <div style="font-size: 2em; font-weight: bold;">1</div> % |
| <hr/> | | |
| <div style="background-color: purple; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">2</div> | <p>bmchealthservres.biomedcentral.com</p> <p>Internet Source</p> | <div style="font-size: 2em; font-weight: bold;">1</div> % |
| <hr/> | | |
| <div style="background-color: purple; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">3</div> | <p>www.mdpi.com</p> <p>Internet Source</p> | <div style="font-size: 2em; font-weight: bold;">1</div> % |
| <hr/> | | |
| <div style="background-color: teal; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">4</div> | <p>pmc.ncbi.nlm.nih.gov</p> <p>Internet Source</p> | <div style="font-size: 2em; font-weight: bold;">1</div> % |
| <hr/> | | |
| <div style="background-color: green; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">5</div> | <p>Zhou Fang, Sijie An, Xixi Sun, Jialu Li, Jiajia Jiao, Yunfei Yin, Ling Wei, Kai Wang, Chengjuan Xie, Xingui Chen, Yubao Jiang. "An Exploratory Study on a Potential Biomarker for Seizures in Autoimmune Encephalitis", Journal of Inflammation Research, 2025</p> <p>Publication</p> | <div style="font-size: 2em; font-weight: bold;">1</div> % |
| <hr/> | | |
| <div style="background-color: brown; color: white; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">6</div> | <p>Yunhao Wang, Huimin Li, Jiewen Wei, Xia Li et al. "Imaging-free hip arthrocentesis with a multipoint infiltration technique for hip pain: a 1-year follow-up pilot real-world study", Journal of Hip Preservation Surgery, 2026</p> <p>Publication</p> | <div style="font-size: 2em; font-weight: bold;"><1</div> % |

7 Yuri Oshiro, Tatsunori Satoh, Emi Ohata, Eiji Nakatani, Hideaki Kaneda, Akira Sugawara. "Development of a Predictive Tool for Maintenance Dialysis Initiation: A Retrospective Cohort Analysis of 589,284 Participants from the Shizuoka Kokuho Database Study", Kidney Medicine, 2025
Publication

8 easts-journal.com
Internet Source

9 research.lstmed.ac.uk
Internet Source

10 Luyun Lin, Yiqing Wang. "SHAP Stability in Credit Risk Management: A Case Study in Credit Card Default Model", Risks, 2025
Publication

11 Submitted to Darius Bwalya
Student Paper

12 www.ncbi.nlm.nih.gov
Internet Source

13 Cankat Genis, Demet Tas, Deniz Yilmaz, Fatma Nur Kuzucu, Ahmet Selmanoglu, Zeynep Sengul Emeksiz, Emine Dibek Mısırlıoglu. ""Unraveling the effects of smoking in asthmatic adolescents: clinical outcomes, spirometry findings, and risk analysis"", European Journal of Pediatrics, 2025
Publication

14 Shorog Althubait, Rawan M. Alqahtani, Razan M. Alqahtani, Wejdan H.A. Al-Qahtani et al. "Public knowledge and awareness of stroke and associated risk factors among adults population in the southern region of Saudi

Arabia: A cross-sectional study", Medicine, 2026

Publication

15

aricjournal.biomedcentral.com

Internet Source

<1 %

16

bmcnutr.biomedcentral.com

Internet Source

<1 %

17

Pinar Uzunkaya Oztoprak, Berna Kurt. "From silence to action: Nursing students' transformative learning experiences in sexuality care for cancer patients – A qualitative study", European Journal of Oncology Nursing, 2026

Publication

<1 %

18

www.primr.org

Internet Source

<1 %

19

Nadeem Aljonaid, Ahmed Alwadeai, Osamah Ahmed AL-Maamari, Mohammed Jaboob, Abdo Al-Barakani, Hadi Mareeh. "Sectoral foreign aid and growth: spatial and temporal dynamics in a regional panel framework", Letters in Spatial and Resource Sciences, 2025

Publication

<1 %

20

dask.kebijakankesehatanindonesia.net

Internet Source

<1 %

21

Shahryar Zeighami, Ali Seifi, Mohammad Hossein Sharifi, Zahra Gheibi et al. "Association of Diabetes and Cancers: An analysis of the Healthcare Cost and Utilization Project National Inpatient Sample (HCUP-NIS) 2005-2015 Dataset", Springer Science and Business Media LLC, 2025

Publication

<1 %

22

bmccancer.biomedcentral.com

Internet Source

<1 %

23

www.archive.aessweb.com

Internet Source

<1 %

24

Ali Kiadaliri, Andrea Dell'Isola, L. Stefan Lohmander, David J. Hunter, Leif E. Dahlberg. "Assessing the importance of predictors of adherence to a digital self-management intervention for osteoarthritis", Journal of Orthopaedic Surgery and Research, 2023

Publication

<1 %

25

Ola Abu Al Karsaneh, Arwa Al Anber, Sahar AlMustafa, Hussien Alma'aitah et al. "Human Papillomavirus Is Rare and Does Not Correlate with p16INK4A Expression in Non-Small-Cell Lung Cancer in a Jordanian Subpopulation", Medicina, 2024

Publication

<1 %

26

Vivian Khamis, Anies Al-Hroub. "School bullying among refugee and non-refugee adolescent students in Lebanon", Child Abuse & Neglect, 2025

Publication

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

Off